


**NASA CONTRACTOR
REPORT**



NASA CR-2314

NASA CR-2314

**THE DISTRIBUTION OF ODD NITROGEN
IN THE LOWER STRATOSPHERE AND
POSSIBLE PERTURBATIONS CAUSED
BY STRATOSPHERIC AIR TRANSPORT**

by Ivar S. A. Isaksen and Eigil Hesstvedt

Prepared by

UNIVERSITY OF OSLO

Oslo, Norway

for Goddard Space Flight Center

NATIONAL AERONAUTICS AND SPACE ADMINISTRATION • WASHINGTON, D. C. • AUGUST 1973

1. Report No. NASA CR-2314	2. Government Accession No.	3. Recipient's Catalog No.	
4. Title and Subtitle THE DISTRIBUTION OF ODD NITROGEN IN THE LOWER STRATOSPHERE AND POSSIBLE PERTURBATIONS CAUSED BY STRATOSPHERIC AIR TRANSPORT		5. Report Date August 1973	
		6. Performing Organization Code	
7. Author(s) IVAR S. A. ISAKSEN AND EIGIL HESSTVEDT		8. Performing Organization Report No.	
9. Performing Organization Name and Address INSTITUTE OF GEOPHYSICS UNIVERSITY OF OSLO OSLO, NORWAY		10. Work Unit No.	
		11. Contract or Grant No. NGR 52-155-002	
12. Sponsoring Agency Name and Address GODDARD SPACE FLIGHT CENTER GREENBELT, MARYLAND 20771 J. THEON, TECHNICAL OFFICER		13. Type of Report and Period Covered FINAL	
		14. Sponsoring Agency Code	
15. Supplementary Notes			
16. Abstract <p>In the lower stratosphere a significant production of odd nitrogen results from the reaction $N_2O + O(^1D) \rightarrow 2NO$. Since the transport is relatively slow, odd nitrogen builds up with a maximum mixing ratio of 2×10^{-8} at 30 km. Profiles of odd nitrogen, for different latitudes, winter and summer, are computed from one-dimensional transport models. Variations with latitude are small. Horizontal transport is therefore not believed to alter our results significantly.</p> <p>In order to evaluate the effect of odd nitrogen upon the ozone layer, NO_x profiles are calculated. OH is here a key component, since it converts NO_2 to HNO_3. In the region where ozone is determined by chemistry rather than by transport (above 25 km), NO_2 is found to be relatively abundant.</p> <p>The effect of stratospheric transport on the NO_x distribution is shown to depend critically upon the height of emission. The effect increases by a factor of 5 or more for a change of flight level from 18 km to 23 km. This strong dependence should be duly considered when future stratospheric transport is discussed.</p>			
17. Key Words (Selected by Author(s)) STRATOSPHERIC COMPOSITION; PHOTO-CHEMISTRY OF ODD NITROGEN IN THE STRATOSPHERE; MODIFICATION OF STRATOSPHERIC COMPOSITION		18. Distribution Statement	
19. Security Classif. (of this report) UNCLASSIFIED	20. Security Classif. (of this page) UNCLASSIFIED	21. No. of Pages 27	22. Price* \$3.00

CONTENTS

	<u>Page</u>
Abstract	1
1. Introduction	1
2. The model	2
3. Atomic oxygen in the excited state and hydroxyl	3
4. Distribution of nitrous oxide	5
5. Production and distribution of odd nitrogen	6
6. The distribution of odd nitrogen on NO, NO ₂ , and HNO ₃	7
7. Air transport in the stratosphere and its effect on the NO _x distribution	9
Acknowledgement	10
References	11

ILLUSTRATIONS

<u>Figure</u>		<u>Page</u>
1	Vertical eddy diffusion coefficient	13
2	Vertical profiles of O (¹ D) for 90° summer, 60° summer, 30° summer, 0°, 30° winter, and 60° winter	14
3	Vertical profiles of OH, for the same latitudes as in Figure 2	15
4	Vertical profiles of N ₂ O for 90° summer, 0°, 30° winter, and 60° winter	16
5	Vertical profiles of N ₂ O loss rates from equations (16) and (18) at 30° summer.	17

ILLUSTRATIONS (continued)

<u>Figure</u>		<u>Page</u>
6	Vertical profiles of production rates of odd nitrogen from equation (16) for the same latitudes as in Figure 4	18
7	Profiles of odd nitrogen mixing ratios for the same latitudes as in Figure 2	19
8	Vertical profiles of odd nitrogen loss rates in the NO - NO ₂ cycle (curves $q_{\text{NO}_2 - \text{NO}}$, and q_{NO}) and in the NO - NO ₂ - HNO ₃ cycle (curves $q_{\text{NO}_2 - \text{HNO}_3}$ and q_{HNO_3}) for 30° summer.	20
9	Distribution of odd nitrogen on NO, NO ₂ , and N ₂ O	21
10	Vertical profiles of NO _x mixing ratio.	22
11	Increase (in %) of NO _x mixing ratio caused by a fleet of 200 SST aircraft as a function of flight level (horizontal axis)	23

THE DISTRIBUTION OF ODD NITROGEN IN THE LOWER STRATOSPHERE AND POSSIBLE PERTURBATIONS CAUSED BY STRATOSPHERIC AIR TRANSPORT

Ivar S. A. Isaksen and Eigil Hesstvedt
Institute of Geophysics
University of Oslo
Oslo, Norway

Abstract


In the lower stratosphere a significant production of odd nitrogen results from the reaction $\text{N}_2\text{O} + \text{O} (^1\text{D}) \rightarrow 2 \text{NO}$. Since the transport is relatively slow, odd nitrogen builds up with a maximum mixing ratio of 2×10^{-8} at 30km. Profiles of odd nitrogen, for different latitudes, winter and summer, are computed from one-dimensional transport models. Variations with latitude are small. Horizontal transport is therefore not believed to alter our results significantly.

In order to evaluate the effect of odd nitrogen upon the ozone layer, NO_x profiles are calculated. OH is here a key component, since it converts NO_2 to HNO_3 . In the region where ozone is determined by chemistry rather than by transport (above 25km), NO_2 is found to be relatively abundant.

The effect of stratospheric transport on the NO_x distribution is shown to depend critically upon the height of emission. The effect increases by a factor of 5 or more for a change of flight level from 18km to 23km. This strong dependence should be duly considered when future stratospheric transport is discussed.

1. Introduction

It is well established (Crutzen, 1971, Johnston, 1971, Hesstvedt, 1973), that nitrogen oxides act upon the ozone layer through the reaction



whereby odd oxygen is destroyed. Accordingly, reliable information about the distribution of nitrogen oxides in the lower stratosphere is essential in quantitative assessments of this effect. Production of odd nitrogen takes place in two regions (Crutzen, 1971, Isaksen, 1973). In the upper region, around 100km,

ionic processes result in an important production of NO and N. In the lower region, between 20 and 30km, NO is produced when N₂O is destroyed by O (¹D). In this region the chemistry appears to be rather complicated. Modeling of the atmosphere with respect to odd nitrogen species, must also comprise a series of oxygen, hydrogen, and carbon species. The model has earlier been presented by Isaksen (1973) and will therefore not be discussed in detail here. We will, however, discuss production and loss reactions with O (¹D) and OH, which are day components for the nitrogen oxides in the lower stratosphere.

2. The model

Important parameters in the NO_x calculations, such as eddy diffusion coefficient, temperature and the air density, are known to vary markedly with season and latitude. Calculations are therefore made for 60° and 30°, winter, for the Equator, and for 30°, 60°, and 90°, summer. Temperatures and densities are taken from CIRA (1965). The models are one-dimensional, and vertical eddy transport is included. The eddy diffusion coefficients used are given in Figure 1. Up to 27km the coefficients are taken from Gudiksen et al., (1968). Above 27km an exponential increase is assumed, as suggested by Lindzen (1971). His indication of a seasonal variation is also adapted.

Hesstvedt (1973) has shown that a two-dimensional model should be considered for computations of the ozone distribution in the lower stratosphere. Since our model is one-dimensional we therefore decided to use observed values of ozone (Hering and Borden, 1964).

Our model is a steady state model, where the photochemistry is averaged over the day. Long lived species which are influenced by eddy diffusion are determined from equations of the type

$$\frac{d [X]}{dt} = - \frac{d}{dz} \left(K_z [M] \frac{d \delta_x}{dz} \right) + P_x - Q_x [X] = 0 \quad (2)$$

where K_z is the vertical eddy diffusion coefficient, M the total number density, $\delta_x = [X] / [M]$ the mixing ratio of component X , and P_x and $Q_x [X]$ are averaged photochemical production and loss terms. This equation is solved by the direct implicit method given by Isaksen (1973), using finite differences.



Components with short chemical lifetimes, hours or less, are not influenced by transport processes. The transport term in equation (2) may then be disregarded, and average daytime values are determined from expressions of the type

$$P_x - Q_x [X] = 0 \quad (3)$$

where P_x and $Q_x [X]$ are average daytime production and loss terms.

3. Atomic oxygen in the excited state and hydroxyl

Two components play an important role in the formation of odd nitrogen species; these components are O (¹D) and OH. Odd nitrogen is produced through the reaction (1) which appears to be the only effective source in the stratosphere. Nitrogen oxides will therefore depend strongly on the O (¹D) profiles, which are determined from




which gives

$$[\text{O} (^1\text{D})] = J_3 \cdot [\text{O}_3] / k_5 \cdot [\text{M}] \quad (6)$$


In our model observed values are used for O₃ and M. J₃ is the daytime average dissociation rate of O₃ for λ < 311 nm. Its strong variation with height in the lower stratosphere (Isaksen, 1973) is a result of ozone absorption, and can therefore be calculated from the observed ozone profile.

O (¹D) profiles, calculated from equation (6) are given in Figure 2 for different latitudes, summer and winter. The rapid decrease below 50km is a result of the decrease in J₃. Variations with latitude are small except at high latitudes, winter, where the production is very slow because of the sun.


Hydroxyl is a second component in the lower stratosphere. It is produced by the reaction






In addition to this reaction there is some production through the reaction



Below 30km, however, this reaction becomes negligible, since O (³P) decreases rapidly below this height. OH is lost through several reactions. Important reactions in the lower stratosphere are found to be






$$f = 1.6 \times 10^{-2} \cdot T^4 / (4 \times 10^{-11} [\text{M}])$$

Reactions (9) and (10) are real loss reactions, since CH_3 and H are ultimately converted to HO_2 . Reaction (12), on the other hand, is not a real loss since most of OH lost by this reaction is reformed through the reaction



The real OH loss is only the fractional part going through reaction (11) and reaction




on the short wavelength side of HNO_3 dissociation. It appears that reaction (14) is much less effective than reaction (13). OH may then be determined from

$$[\text{OH}] = \frac{(k_7 [\text{NO}] + k_8 [\text{O}(^3\text{P})]) [\text{HO}_2]}{k_9 [\text{CH}_4] + K_{10} [\text{CO}] + 2k_{11} [\text{HNO}_3] + J_{\text{bHNO}_3}} \quad (15)$$


OH can therefore only be determined in a model where all the components occurring in equation (15) are computed simultaneously. This is done in our model. Calculations are repeated until convergence is obtained. The resulting OH profiles are shown in Figure 3. OH has, for all latitudes, a maximum near 45km with a peak value of $2 - 3 \times 10^{17} \text{ cm}^{-3}$. Down to the peak level HO_2 is effectively converted to OH through reaction (8), while the destruction through reactions (9), (10), and (12) is slow. Below 45km this is no longer true and we get a rapid decrease in OH down to the tropopause, where we find a minimum value near 10^5 cm^{-3} . The strong variations with latitude below about 17km come from the variations in the tropopause level. At the Equator heights below 17km are in the troposphere, at high latitudes all heights above 8km are in the stratosphere.

4. Distribution of nitrous oxide


Destruction of N_2O through the reaction



is the only important source of NO_x in the lower stratosphere. Realistic information on the spatial distribution of this component is therefore essential. N_2O has been found to be fairly uniformly distributed in the troposphere with negligible seasonal variations. A mixing ratio of $2.5 \cdot 10^{-7}$ is regarded as representative for the troposphere (Schütz et al., 1970). Production of N_2O is expected to be through soil bacteria. So far no photochemical sources are known. It is therefore difficult to estimate production rates. Since the loss rate is small, vertical transport maintain a constant mixing ratio of N_2O in the troposphere, and the above value is used as boundary value at the tropopause. In the lower stratosphere the conditions are quite different. Diffusion is much lower and loss becomes more effective through the reaction



and through the slower reaction



in addition to reaction (16). N_2O profiles are therefore obtained from

$$-\frac{d}{dz} \left(K_z [M] \frac{d}{dz} (\delta_{N_2O}) \right) - (J_{N_2O} + (k_{16} + k_{18}) [O(^1D)]) [N_2O] = 0 \quad (19)$$

The results are shown in Figure 4. The slope of the profiles depends on the dissociation rates and on the eddy diffusion coefficient. Weak diffusion leads to a rapid decrease in the N_2O mixing ratio (δ_{N_2O}). This occurs at 25-30km, where K_z has a minimum. Variations with latitude and season are relatively small, except for high latitude, winter. Apart from this case, production of NO_x through destruction of N_2O is confined to levels below 35-40km. At 60°, winter, the low sun makes photodissociation less important, and when in addition the eddy mixing is stronger, N_2O is transported to greater heights before it is dissociated.

5. Production and distribution of odd nitrogen


From the profiles of N_2O and $O(^1D)$, shown in Figures 2 and 4, the production rates of odd nitrogen are easily calculated from

$$P_{NO_y} = 2k_{16} \cdot [O(^1D)] [N_2O] \quad (20)$$

The efficiency of this production depends on how efficient reaction (16) is in breaking down N_2O , compared to reactions (17) and (18). Figure 5 shows the loss rate of N_2O from reactions (16) and (17). Loss by reaction (18) is equal to the loss by reaction (16). The curves are for 30° summer. It is clearly demonstrated that reaction (17) is the main loss reaction of N_2O . Only a few percent of the loss will produce nitrogen oxides through reaction (16).

Nitrogen oxide production from eq. (20) is given in Figure 6. In the summer production varies relatively little with latitude. Differences below 20km is a result of the variation in tropopause level, which influences the $O(^1D)$ densities. At all latitudes, odd nitrogen production increases rapidly up to a marked maximum between 22-26km. In the winter the production is important over a much wider region. This is clearly demonstrated for 60°, winter, where a substantial production takes place above 40km, with a maximum around 35km. The winter profiles are a result of the combined effects of the absorption of radiation producing $O(^1D)$ and the strong eddy diffusion, which effectively transports N_2O to greater heights. At 30°, winter, these effects are less pronounced, with the result that we get a profile more like the summer profiles. Maximum production rates vary from $1.5 \times 10^2 \text{ cm}^{-3} \text{ s}^{-1}$ in the summer hemisphere, to $1.5 \times 10^1 \text{ cm}^{-3} \text{ s}^{-1}$ at 60°, winter. If we add up the total production in a column, the average production for the whole globe becomes $10^8 \text{ cm}^{-2} \text{ s}^{-1}$.

The distribution of total odd nitrogen in the lower stratosphere is easily calculated when its production rate is known. The only photochemical loss of nitrogen oxides in the atmosphere is through the reaction



However, atomic nitrogen decreases rapidly below 60km (Isaksen, 1973). This reaction is therefore of no interest for our problem. Odd nitrogen is therefore removed from the lower stratosphere by vertical eddy diffusion alone. Since K_z increases with height, the particle flux of odd nitrogen, NO_y

$$F = -K_z [M] \frac{d}{dz} (\delta_{NO_y}) \quad (22)$$

is directed downwards.

The flux is proportional to the eddy diffusion coefficient and to $[M]$. It is, therefore clear that the increase in δ_{NO_y} , sufficient to transport away the odd nitrogen produced at a certain height, depends strongly on the heights where the production takes place. Assuming steady state conditions,


$$-\frac{d}{dz} \left(K_z [M] \frac{d}{dz} (\delta_{NO_y}) \right) + P_{NO_y} = 0 \quad (23)$$

gives the height distribution of nitrogen in all odd forms, i.e., in the forms of NO, NO₂, and HNO₃; all other odd components are present in negligible quantities. Results of this calculation are given in Figure 7. As might be expected from the considerations above, there are latitudinal variations in odd nitrogen, due to variations in K_z and in production rates. Above 20km, where the effect on ozone is of importance, variations with latitude are very small in the summer. The mixing ratio of odd nitrogen increases strongly up to 25km, the region of the strongest production (Fig. 6). But even above this height there is substantial increase to above 30km, partly due to the low K_z values. Above 35km δ_{NO_y} drops as a result of the low mesosphere values used for NO (Isaksen, 1973 a). Maximum mixing ratios of 1.2×10^{-8} are found around 32km.


In the winter the low solar elevation leads to production at higher altitudes. Therefore the increase in mixing ratio takes place at higher altitudes. This is especially true for 60°, winter, where the strong atmospheric absorption gives high mixing ratios in the upper stratosphere. It is, therefore, an increase in odd nitrogen mixing ratios in the whole stratosphere.

6. The distribution of odd nitrogen on NO, NO₂, and HNO₃

In order to estimate the effect on ozone, we shall next see how the total odd oxygen, as computed above, is distributed on the different odd nitrogen species. The initial product is NO through reaction (16). This component will rapidly be converted to NO₂ through reaction




The balance between NO and NO₂ is maintained through solar dissociation of NO₂



with some contribution from reaction (1).

The conversion to HNO_3 is much slower (reaction (12)). To obtain HNO_3 densities, the loss reactions must be compared with reaction (12). The main loss of HNO_3 is through reactions (13) and (14) with some contribution from reaction (11). NO_3 is rapidly converted to NO and NO_2 through the reactions



NO_3 is unlikely to be of importance and will be disregarded here.

From the above reactions the distribution of odd nitrogen on NO , NO_2 , and HNO_3 can be calculated. Equilibrium between NO and NO_2 is given by reactions (24), (25), and (1). Loss rates of NO and NO_2 are shown in Figure 8 for 30° summer. It is seen that equilibrium is established after a few minutes. Similarly, equilibrium between $\text{NO} - \text{NO}_2$ and HNO_3 is given by equations (11) - (14). Equilibrium in this cycle is much slower, as shown in the same figure. Lifetimes decrease from 1 day at 30km to 10 days at 15km. The short lifetimes in the NO to NO_2 cycle means that NO is converted to NO_2 a few minutes after sunset, when there is no photodissociation. Likewise, NO is also reproduced rapidly after sunrise from photodissociation of NO_2 .

The long lifetimes in the $\text{NO} - \text{NO}_2 - \text{HNO}_3$ cycle result in negligible HNO_3 variations over the day below 30km. By confining the equations for equilibrium in the $\text{NO} - \text{NO}_2$ and in the $\text{NO} - \text{NO}_2 - \text{HNO}_3$ cycle, average daytime number densities for NO , NO_2 , and HNO_3 are given by

$$[\text{NO}] = [\text{NO}_y] \frac{J_{\text{HNO}_3} + k_{11} [\text{OH}]}{J_{\text{HNO}_3} + k_{11} [\text{OH}] + k'_{12} [\text{OH}] [\text{M}]} \cdot \frac{k_1 [\text{O} (^3\text{P})] + J_{\text{NO}_2}}{k_1 [\text{O}] + J_{\text{NO}_2} + k_{24} [\text{O}_3]}\tag{27}$$

$$[\text{NO}_2] = [\text{NO}_y] \frac{J_{\text{HNO}_3} + k_{11} [\text{OH}]}{J_{\text{HNO}_3} + k_{11} [\text{OH}] + k'_{12} [\text{OH}] [\text{M}]} \cdot \frac{k_{24} [\text{O}_3]}{k_1 [\text{O} (^3\text{P})] + J_{\text{NO}_2} + k_{24} [\text{O}_3]}\tag{28}$$

$$[\text{HNO}_3] = [\text{NO}_y] \frac{k'_{12} [\text{OH}] [\text{M}]}{J_{\text{HNO}_3} + k_{11} [\text{OH}] + k'_{12} [\text{OH}] [\text{M}]} \quad (29)$$

k'_{12} gives the efficiency of the coefficient k_{12} in converting $\text{NO} + \text{NO}_2$ to HNO_3 :

$$k'_{12} = k_{12} \frac{k_{24} [\text{O}_3]}{k_{24} [\text{O}_3] + k_1 \cdot [\text{O} (^3\text{P})] + J_{\text{NO}_2}} \quad (30)$$

The relative abundances of the three nitrogen species, given by equations (27), (28), and (29), are presented in Figure 9 for heights between 10km and 40km for 30° summer. HNO_3 is seen to be the dominating component below about 27km. At 25km 70% of odd nitrogen present is in the form of HNO_3 , and it exceeds 40% at all heights below. The maximum HNO_3 concentrations around 25km is a result of the high ozone concentration. Below about 15km HNO_3 increases due to the increase in OH. Above 35km HNO_3 is present in small quantities. NO_2 , which is of special interest in the ozone chemistry, is most abundant around 30km, where 50% of the total odd nitrogen is in the form of NO_2 . Relative densities vary little below 25km, between 20% and 30%. NO is the most abundant component above 35km, compared with the other components. Below 30km it is a minor component. The NO - NO_2 cycle is of interest for ozone destruction through reaction (1). As we have already seen, lifetimes are very short. During the day, when ozone is lost through reaction (1), NO and NO_2 are in photochemical equilibrium. When the sum of NO and NO_2 is known, equilibrium values are easily calculated from equations (1), (24), and (25).

NO_x mixing ratios are given in Figure 10 for summer and winter hemisphere. Above 30km the mixing ratios differ little from odd nitrogen mixing ratios, due to low HNO_3 densities (Fig. 8). Below 25km, there is a sharp drop down to 10km, where mixing ratios of about 10^{-9} are obtained for all latitudes. Variations with latitude are very small, except in the winter hemisphere where δ_{NO_x} falls off at higher latitude.

7. Air transport in the stratosphere and its effect on the NO_x distribution

It is evident that nitrogen oxides produced from aircraft flying in the stratosphere will alter the stratospheric NO_x distribution. Since vertical eddy diffusion varies strongly with height, it is of interest to estimate the magnitude of this additional source when the aircraft are assumed to fly at given heights. We shall here consider the effect of a "reference fleet" of 200 four-engine aircraft, each performing 1000 flights of 2-1/2 hours per year. The fuel flow is assumed

to be 5000kg per hour per engine. The emission of NO (converted to NO₂) is assumed to be 21 g per kg fuel. These 200 aircraft are assumed to fly at the same flight level uniformly routed around the globe. This gives an average production rate of 1.88×10^7 NO_x molecules cm⁻²s⁻¹. Since we use finite differences with steps of 1km, we have, in our model, a production rate of 1.88×10^2 NO_x molecules cm⁻²s⁻¹. This number will be used as an additional source in equation (23) at given flight levels. The variation in δ_{NO_x} , for flight levels between 15km and 32km, is given in Figure 11. The steady state profiles in Figure 10 for 60°, summer, and 30°, winter, are used as a basis, and the increase is given in percent. Since the ozone distribution below 25km is largely dominated by transport processes (Hesstvedt, 1973 a), the increase in NO_x between 25km and 30km is of primary interest for the ozone problem.

As might be expected, the increase in NO_x depends strongly on the flight level. NO_x injection at high altitudes leads to a considerable increase in NO_x. This can be explained by the very fast drop in eddy diffusion coefficient above the tropopause (Fig. 1). The added NO_x has to be transported downwards into the troposphere. Production at heights where K_z is small, therefore means that the gradient in δ_{NO_x} must be much greater than the gradient is at lower heights with higher K_z values, in order to maintain the flux of NO_x given by equation (22).

The increase in δ_{NO_x} is somewhat different in the two diagrams of Figure 11, a result of difference in K_z . In both cases, however, the perturbation in NO_x increases rapidly as the source level, given along the abscissa, is moved upwards. With the source at 18km, the increase in NO_x is approximately 2% for 60°, summer, and 0.5% for 30°, winter. With the source at 23km, NO_x increases with 5-10% in the 25-30km region in both models, and with source at 28km, NO_x increases by around 20% in both models. The calculations give similar results for the other latitudes. These results have been used in a two-dimensional model to show the effect on the ozone layer (Hesstvedt, 1973 a).

These numbers clearly demonstrate that the effect of stratospheric transport on the NO_x densities depends strongly on the flight levels. Flight levels above 20km lead to a considerable increase in NO_x. Having in mind that the perturbation (in %) of the ozone layer is roughly one third of the perturbation (in %) of NO_x, the importance of the flight level should be duly considered when the effect of stratospheric transport upon the ozone layer is discussed.

Acknowledgement

This work was partially supported by National Aeronautics and Space Administration grant NGR 52-155-002.

References

1. CIRA. 1965. International Reference Atmosphere. North Holland Publ. Co., Amsterdam, 1965.
2. Crutzen, P. J., 1971. J. Geophys. Res., 76, 7311-7327.
3. Gudiksen, P. H., Fairhall, A. W., & Reed, R. J., 1968. J. Geophys. Res., 77, 4461-4473.
4. Hering, W. S., & Borden, T. S., 1964. Vol. 2. Environmental Research Papers, No. 38.
5. Hesstvedt, E., 1973. Water, Air and Soil Pollution (in print).
6. Hesstvedt, E., 1973 a. Paper presented at the AGARD Symposium in London, 9-13 April 1973, (in print).
7. Isaksen, I. S. A., 1973. Geophysica Norvegica (in print).
8. Isaksen, I. S. A., 1973 a. Nitric oxide distribution in the mesosphere. To be published.
9. Johnston, H. F., 1971. Science 173, 511-522.
10. Lindzen, R. S., 1971. In Mesospheric Models and Related Experiments. Reidel, 122-130.
11. Schütz, K., Junge, C., Beck, R., & Albrecht, B., 1970. J. Geophys. Res., 75, 2230-2246.

Figure Captions

- Figure 1. Vertical eddy diffusion coefficient.
- Figure 2. Vertical profiles of O (^1D) for 90° summer, 60° summer, 30° summer, 0°, 30° winter, and 60° winter.
- Figure 3. Vertical profiles of OH, for the same latitudes as in Figure 2.
- Figure 4. Vertical profiles of N_2O for 90° summer, 0°, 30° winter, and 60° winter.
- Figure 5. Vertical profiles of N_2O loss rates from equations (16) and (18) at 30° summer.
- Figure 6. Vertical profiles of production rates of odd nitrogen from equation (16) for the same latitudes as in Figure 4.
- Figure 7. Profiles of odd nitrogen mixing ratios for the same latitudes as in Figure 2.
- Figure 8. Vertical profiles of odd nitrogen loss rates in the $\text{NO} - \text{NO}_2$ cycle (curves $q_{\text{NO}_2 - \text{NO}}$ and q_{NO}) and in the $\text{NO} - \text{NO}_2 - \text{HNO}_3$ cycle (curves $q_{\text{NO}_2 - \text{HNO}_3}$ and q_{HNO_3}) for 30° summer.
- Figure 9. Distribution of odd nitrogen on NO , NO_2 , and HNO_3 .
- Figure 10. Vertical profiles of NO_x mixing ratio.
- Figure 11. Increase (in %) of NO_x mixing ratio caused by a fleet of 200 SST aircraft as a function of flight level (horizontal axis).




Figure 1. Vertical eddy diffusion coefficient.




Figure 2. Vertical profiles of O(1D) for 90° summer, 60° summer, 30° summer, 0° summer, 0° winter, 30° winter, and 60° winter.




Figure 3. Vertical profiles of OH, for the same latitudes as in Figure 2.




Figure 4. Vertical profiles of N_2O for 90° summer, 0° summer, 30° winter, and 60° winter.




Figure 5. Vertical profiles of N_2O loss rates from equations (16) and (18) at 30° summer.




Figure 6. Vertical profiles of production rates of odd nitrogen from equation (16) for the same latitudes as in Figure 4.




Figure 7. Profiles of odd nitrogen mixing ratios for the same latitudes as in Figure 2.




Figure 8. Vertical profiles of odd nitrogen loss rates in the NO-NO₂ cycle (curves q_{NO_2-NO} and q_{NO}) and in the NO-NO₂-HNO₃ cycle (curves $q_{NO_2-HNO_3}$ and q_{HNO_3}) for 30° summer.




Figure 9. Distribution of odd nitrogen on NO , NO_2 , and HNO_3 .




Figure 10. Vertical profiles of NO_x mixing ratio.




Figure 11. Increase (in %) of NO_x mixing ratio caused by a fleet of 200 SST aircraft as a function of flight level (horizontal axis).



POSTMASTER : If Undeliverable (Section 158
Postal Manual) Do Not Return

"The aeronautical and space activities of the United States shall be conducted so as to contribute . . . to the expansion of human knowledge of phenomena in the atmosphere and space. The Administration shall provide for the widest practicable and appropriate dissemination of information concerning its activities and the results thereof."

—NATIONAL AERONAUTICS AND SPACE ACT OF 1958

NASA SCIENTIFIC AND TECHNICAL PUBLICATIONS

TECHNICAL REPORTS: Scientific and technical information considered important, complete, and a lasting contribution to existing knowledge.

TECHNICAL NOTES: Information less broad in scope but nevertheless of importance as a contribution to existing knowledge.

TECHNICAL MEMORANDUMS: Information receiving limited distribution because of preliminary data, security classification, or other reasons. Also includes conference proceedings with either limited or unlimited distribution.

CONTRACTOR REPORTS: Scientific and technical information generated under a NASA contract or grant and considered an important contribution to existing knowledge.

TECHNICAL TRANSLATIONS: Information published in a foreign language considered to merit NASA distribution in English.

SPECIAL PUBLICATIONS: Information derived from or of value to NASA activities. Publications include final reports of major projects, monographs, data compilations, handbooks, sourcebooks, and special bibliographies.

TECHNOLOGY UTILIZATION PUBLICATIONS: Information on technology used by NASA that may be of particular interest in commercial and other non-aerospace applications. Publications include Tech Briefs, Technology Utilization Reports and Technology Surveys.

Details on the availability of these publications may be obtained from:

SCIENTIFIC AND TECHNICAL INFORMATION OFFICE

NATIONAL AERONAUTICS AND SPACE ADMINISTRATION
Washington, D.C. 20546